

University of Groningen

## SOY MILK FERMENTATION

De Bok Franciscus Adrianus, Maria; Janssen Patrick Wilhelmus, Maria; Sikkema, Jan; Smid Eilt, Johannes

**IMPORTANT NOTE:** You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

*Publication date:*  
2012

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

De Bok Franciscus Adrianus, M., Janssen Patrick Wilhelmus, M., Sikkema, J., & Smid Eilt, J. (2012). SOY MILK FERMENTATION. (Patent No. *US2012177782*).

### Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

### Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

*Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.*



US 20120177782A1

(19) **United States**(12) **Patent Application Publication**  
**De Bok et al.**(10) **Pub. No.: US 2012/0177782 A1**(43) **Pub. Date: Jul. 12, 2012**(54) **SOY MILK FERMENTATION****Related U.S. Application Data**(75) Inventors: **Franciscus Adrianus Maria De Bok**, Wageningen (NL); **Patrick Wilhelmus Maria Janssen**, Ede (NL); **Jan Sikkema**, Zeegse (NL); **Eilt Johannes Smid**, Wageningen (NL)

(60) Provisional application No. 61/241,117, filed on Sep. 10, 2009.

(30) **Foreign Application Priority Data**

Sep. 10, 2009 (EP) ..... 09169906.6

**Publication Classification**(73) Assignee: **DSM IP ASSETS B.V.**, Heerlen (NL)(51) **Int. Cl.**  
**A23L 1/20** (2006.01)  
**C12N 1/20** (2006.01)  
**C12Q 1/02** (2006.01)(21) Appl. No.: **13/394,772**(52) **U.S. Cl.** ..... **426/46**; 435/29; 435/252.9; 426/61(22) PCT Filed: **Sep. 10, 2010**(57) **ABSTRACT**(86) PCT No.: **PCT/NL2010/050572**§ 371 (c)(1),  
(2), (4) Date: **Mar. 19, 2012**

The present invention relates to a method for preparing a soy-based fermented food product using *L. delbrueckii* and a strain capable of stimulating growth of *L. delbrueckii*, and a food product obtainable by such method. The invention also provides for a culture comprising said strains, and the use thereof for preparing said fermented food product.

Fig 1a





Fig 1b



## SOY MILK FERMENTATION

### FIELD OF THE INVENTION

[0001] The present invention is in the field of fermentation of an aqueous medium comprising soy protein as main protein constituent, in particular to obtain soy yogurt. The invention provides for a method for production of a soy-based fermented food product, and a soy-based fermented food product thus obtained. The invention also relates to new cultures for the preparation of such soy-based fermented food product.

### BACKGROUND

[0002] According to the Codex Alimentarius (FAO/WHO 1977), yogurt is milk fermented with symbiotic cultures of *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus*. Also, *Lactobacillus delbrueckii* subsp. *lactis* is sometimes used in the preparation of dairy yogurt.

[0003] In soy yogurts the main source of protein is not milk, but soy. Soy yogurts are usually fermented with yogurt culture, i.e., symbiotic cultures of *Lactobacillus bulgaricus* (*Lactobacillus delbrueckii* subsp. *bulgaricus*) and *Streptococcus thermophilus*, optionally in conjunction with further lactic acid bacteria.

[0004] Soybeans are less expensive and more abundant than bovine milk, and do not contain cholesterol. Yogurt is one of the fastest growing cultured dairy product in the Western world. Soy yogurt may command a market on the strength of being cholesterol-free. Thus, the economic incentive exists to develop a soy yogurt, in which soy can be utilized as a major ingredient. The basis for such soy yogurt may be soy milk, originating from the soybean. The major carbohydrates in soybeans are sucrose, raffinose and stachyose. The main problem limiting widespread consumption of soybeans and derivatives thereof such as soy yogurt is their objectionable beany flavor and high levels of the oligosaccharides raffinose and stachyose, which can lead to flatulence when consumed in considerable amounts.

[0005] *Lactobacillus delbrueckii* subsp. *bulgaricus* (hereinafter also referred to as "*Lactobacillus delbrueckii bulgaricus*", "*Lactobacillus bulgaricus*" or "*L. bulgaricus*") generally shows limited outgrowth in soy milk compared to bovine milk. This affects the properties of a fermented soy product which has been fermented using *Lactobacillus delbrueckii* subsp. *bulgaricus*. The present inventors have now found that certain *Lactobacillus plantarum* strains improve outgrowth of *Lactobacillus delbrueckii* subsp. *bulgaricus* in soy protein-comprising media, thereby improving the yogurt-type product that is obtained by fermenting soy milk using *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Streptococcus thermophilus* compared to such product obtained by fermenting in the absence of such *Lactobacillus plantarum* strains.

[0006] *L. delbrueckii* subsp. *bulgaricus* degrades sucrose, which is one of the main sugar constituents present in soy milk, poorly. Without wishing to be bound by theory, it is hypothesized that the *Lactobacillus plantarum* strains affecting growth of *L. delbrueckii* subsp. *bulgaricus* are capable of breaking down sucrose, raffinose and/or stachyose molecules into substrates allowing growth of *L. delbrueckii* subsp. *bulgaricus*.

### SUMMARY OF THE INVENTION

[0007] A combinatorial screening approach to identify strain-dependent traits related to mixed culture performance

in natural substrates was developed and used to screen a collection of *L. plantarum* strains for 2- and 3-species mixed culture growth with *Streptococcus thermophilus* CNRZ1066 and *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC-BAA365 in milk and soymilk. This revealed a strain dependent stimulatory effect of *L. plantarum* B1839, B2484, and B2485 on *L. bulgaricus* in soymilk, resulting in a mutualistic relationship independent of the presence of *S. thermophilus*. While in 38 out of 40 mixed cultures with *L. plantarum*, the levels of *L. bulgaricus* were below the detection limit (500 CFU/ml) after 4 rounds of fermentation in soymilk (5-ml static batch fermentations for 24 h at 37° C.), strain B1839 stimulated *L. bulgaricus* over a 100-fold when compared to the corresponding monoculture, yielding nearly 10<sup>8</sup> CFU/ml. The effect was also observed in the 3-species mixed cultures further including *S. thermophilus* yielding a 3-species stable culture. The stimulatory effect could be reproduced in 1 L stirred batch fermentations which were performed to study the population dynamics of all possible combinations of *L. plantarum* strain B1839 in soymilk. Clearly strain B1839 stimulated growth of *L. bulgaricus* in natural soymilk (no additional sugars added) and was thereby able to confer yoghurt-like properties to fermented soymilk products. In 3-species mixed cultures, also strains B2484 and B2485 had a stimulatory effect on *L. bulgaricus* in soy milk.

[0008] Thus, in a first aspect the present invention provides for a method for preparing a fermented soy product, said method comprising the step of fermenting an aqueous medium comprising 0.25-10% (w/w) soy protein with a mixed bacterial culture comprising, in addition to *Lactobacillus delbrueckii*, a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

[0009] In a second aspect, the invention pertains to a fermented soy product comprising *Lactobacillus delbrueckii* and a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

[0010] In a further aspect, the invention relates to a culture comprising *Lactobacillus delbrueckii* and a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

[0011] In other aspects, the invention provides for the use of *Lactobacillus plantarum* B1839, B2482 and/or B2485 to stimulate growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC-BAA365, ATCC-11842 or 1b, in an aqueous medium comprising 0.25-10% (w/w) soy protein, as well as its use in mixed culture fermentation of an aqueous medium comprising 0.25-10% (w/w) soy protein.

[0012] In yet another aspect, the invention sees to a method for identifying a lactic acid bacterial strain capable of stimulating growth of *Lactobacillus delbrueckii* in a medium, said method comprising the steps of: a) growing a monoculture of *Lactobacillus delbrueckii* in said medium; b) growing a mixed culture of *Lactobacillus delbrueckii* and said lactic acid bacterial strain in said medium; c) comparing the growth of *Lactobacillus delbrueckii* in said monoculture with that in said mixed culture; and d) selecting the lactic acid bacterial strain which increases colony forming units *Lactobacillus delbrueckii* in said medium.

### DETAILED DESCRIPTION OF EMBODIMENTS

[0013] Screening of the *L. plantarum* strains for mixed culture fermentation properties in soymilk yielded 40 *L. plantarum*-*S. thermophilus* cultures with high CFU levels of both

species. This combination also dominated the 3-species mixed cultures including *L. bulgaricus*, however, *L. bulgaricus* was still detected in 5 of these cultures. Since *L. bulgaricus* was detected in all monoculture controls as well in the mixed culture controls with *S. thermophilus* it was concluded that growth of this species in mixed cultures with *L. plantarum* as well as in the 3-species cultures was *L. plantarum* strain dependent. Three of the *L. plantarum* strains, B1839, B2484, and B2485, even resulted in increased CFU levels of *L. bulgaricus* in the corresponding 3-species combinations. With *L. plantarum* strain B1839 the *L. bulgaricus* CFU level was highest: 171- and 23-fold higher than in the corresponding mono- and mixed culture controls, respectively. For this reason, strain B1839 was selected to study mono- and mixed-culture population dynamics during soymilk fermentation. In a batch fermenter *L. bulgaricus* levels remained below the detection limit (of 500 CFU/ml), but *L. bulgaricus* grew well in each of the mixed cultures. Also in the batch fermenter a substantial positive effect of *L. plantarum* B1839 and *L. bulgaricus* on each other was observed. Most likely the positive effect of *L. bulgaricus* on *L. plantarum* was related to proteolytic activity since strain B1839 was not able to degrade the available sucrose completely in the monoculture. Since *L. bulgaricus* only poorly ferments sucrose, we hypothesize that the stimulatory effect of *L. plantarum* on *L. bulgaricus* is related to release of sugar monomers from the soymilk sugars by *L. plantarum*.

[0014] In a first aspect, the present invention pertains to a method for preparing a fermented soy product, said method comprising the step of fermenting an aqueous medium comprising 0.25-10% (w/w) soy protein with a mixed bacterial culture comprising, in addition to *Lactobacillus delbrueckii*, a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

[0015] The term "*Lactobacillus delbrueckii*" as used herein includes the subspecies *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Lactobacillus delbrueckii* subsp. *lactis*.

[0016] The amount of soy protein in the fermented soy product according to the invention is 0.25-10% (w/w), preferably 0.5-10% (w/w), such as 0.5-8% (w/w), 0.5-7% (w/w), 1-7% (w/w), 2-6% (w/w), 3-6% (w/w), 4-6% (w/w). It is prepared using an aqueous medium comprising 0.25-10% (w/w), preferably 0.5-10% (w/w), 0.5-8% (w/w), such as 0.5-7% (w/w), 1-7% (w/w), 2-6% (w/w), 3-6% (w/w), 4-6% (w/w) soy protein. For example, such aqueous medium may be soy milk. Soy milk is a water extract of soy beans. Soy milk is generally made by grinding and heating soy beans, removing the fibrous okara (soy pulp), clarifying, and pasteurizing into a soy base. The term "soy protein" as used herein includes soy protein isolate, which may be (partially) hydrolysed, and soy protein concentrate, which may be (partially) hydrolysed.

[0017] Liquid soymilk, soy milk powder, soy protein isolate and soy protein concentrate, and the like are commercially available, e.g. from Devansoy.

[0018] A mixed bacterial culture as used in the present invention comprises at least one *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, strain as well as at least one strain, preferably a lactic acid bacterial strain, capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*. As used herein, the term "lactic acid bacteria" refers to gram-positive rod- or sphere-shaped bacteria that produce lactic acid as the principal or sole end product of

carbohydrate fermentation. Well-known genera include *Bifidobacterium*, *Carnobacterium*, *Enterococcus*, *Lactobacillus*, *Lactococcus*, *Lactospaera*, *Leuconostoc*, *Oenococcus*, *Pediococcus*, *Streptococcus*, *Vagococcus* and *Weissella*. Said mixed bacterial culture may further comprise additional lactic acid bacteria, such as *Streptococcus thermophilus*, *Lactobacillus acidophilus*, *Lactobacillus* subsp. *casei*. In an embodiment, said mixed bacterial culture further comprises at least *Streptococcus thermophilus*. The mixed bacterial culture may also comprise non-lactic acid bacteria, e.g., bacteria of the genus *Propionibacterium*, or any other food grade bacteria.

[0019] Said *Lactobacillus delbrueckii* subsp. *bulgaricus* may be any *Lactobacillus delbrueckii* subsp. *bulgaricus* strain. The skilled person is capable of selecting *Lactobacillus delbrueckii* subsp. *bulgaricus* strains that may be used in the preparation of the fermented soy product of the present invention.

[0020] The fermentation of the aqueous medium comprising 0.25-10% (w/w) soy protein may include one or more of the following steps: a) providing said aqueous medium; b) subjecting said aqueous medium to thermal treatment and homogenisation; c) addition of said mixed bacterial culture; and d) allowing fermentation to take place at a temperature in the range of 33-45° C., preferably 35-43° C., even more preferably 37-40° C.

[0021] In an embodiment, said strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus* stimulates growth of *L. delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in a sucrose-comprising aqueous medium, said medium preferably comprising sucrose and the oligosaccharides stachyose and raffinose as sole carbohydrates in addition to the 0.5-10% (w/w) soy protein. Thus, in a suitable embodiment, said aqueous medium further comprises sucrose, stachyose and/or raffinose as sole carbohydrates. Preferably, said sucrose is present in said aqueous medium in an amount of about 0.25 to about 20% (w/w), preferably 0.25-18% (w/w), 0.4-17% (w/w), 0.5-15% (w/w), 1-13% (w/w). "(w/w)", as herein used, is an abbreviation for "by weight," used to describe the concentration of a substance in a mixture or solution. As an example, a weight percentage of 2, 2% (w/w), means that the mass of the substance is 2% of the total mass of the aqueous medium, i.e. 2 g per 100 g aqueous medium.

[0022] In a suitable embodiment, said strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, is a *Lactobacillus plantarum* strain. Said *Lactobacillus plantarum* strain may be one or more strain selected from the group consisting of *Lactobacillus plantarum* B1839, B2484 and B2485. The *Lactobacillus plantarum* strain B1839 referred to herein corresponds to *Lactobacillus plantarum* SF2A35B published by Figuero et al. (1995. Lactic acid bacteria of the sour cassava starch fermentation. Lett. Appl. Microbiol. 21:126-130). It has been deposited at the Centraalbureau for Schimmelcultures under the Regulations of the Budapest Treaty, and received the deposit number CBS 125104. *Lactobacillus plantarum* strains B2484 (NCTH19-1) and B2485 (NCTH19-2) have been derived from sour pickled pork sausage from Vietnam (Nem Chua). "Nem chua" is made from minced pork, sliced pigskin and a mixture of powdered grilled rice, salt, pepper, sugar and garlic. These contents are mixed thoroughly before being wrapped with aromatic, fresh

leaves into small, boxy rolls before being stored for the fermentation process for three to five days, and when it is ready it can be served direct or grilled on the charcoal stove. B2484 and B2485 have been isolated after the fermentation process, and prior to further processing. They have been deposited at the Centraalbureau for Schimmelcultures under the Regulations of the Budapest Treaty, and received the deposit numbers CBS 125105 (B2484) and CBS 125106 (B2485).

[0023] In a preferred embodiment, said strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, is *Lactobacillus plantarum* B1839, as the stimulatory effect on *Lactobacillus delbrueckii* subsp. *bulgaricus* in an aqueous medium comprising 0.25-10% (w/w) soy protein was most pronounced for this strain.

[0024] In an embodiment, said *Lactobacillus delbrueckii* subsp. *bulgaricus* is selected from the group consisting of *Lactobacillus delbrueckii* *bulgaricus* ATCC-BAA365, ATCC-11842, and Ib.

[0025] In a suitable embodiment, the fermented soy product is soy yogurt. The soy yogurt of the invention may be set-style yogurt with a firm gelled texture, or may be stirred-style yogurt with a spoonable or fluid texture. In an embodiment, the fermentation for the set-style soy yogurt takes place in the package of the yogurt after packaging. Stirred-type soy yogurt is preferably fermented in fermentation tanks. After fermentation, the product may be stirred at a low speed prior to its transport by pumping and filling. A moderate to low shear rate is critical to achieve the desired viscous properties of stirred-style yogurt. Sweeteners, salt and flavors, calcium, fruit and vitamins A and D may be added to the soy yogurt. Stabilizers may also be used in soy yogurt to improve the body and texture by increasing firmness, and, if fruit is included in the soy yogurt, helping to keep the fruit uniformly mixed in the soy yogurt. Stabilizers conventionally used in soy yogurt include alginates (carageenan), gelatins, gums (locust bean, guar), pectins, and starch. Further conventional soy yogurt ingredients may be used. One skilled in the art will be capable of determining suitable conventional soy yogurt ingredients.

[0026] In a further aspect, the present invention pertains to a fermented soy product comprising *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, and a strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*.

[0027] Preferably, said strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in particular in an aqueous medium comprising 0.25-10% (w/w) soy protein, is a *Lactobacillus plantarum* strain, preferably selected from the group consisting of *Lactobacillus plantarum* B1839, B2484, and B2485, more preferably *Lactobacillus plantarum* B1839.

[0028] In an embodiment, said *Lactobacillus delbrueckii* *bulgaricus* is selected from the group consisting of *Lactobacillus delbrueckii* *bulgaricus* ATCC-BAA365, ATCC-11842, and Ib. As outlined hereinabove, in an advantageous embodiment, the fermented soy product is soy yogurt.

[0029] The fermented soy product may comprise further food ingredients such as sugars, further proteins, fats and oils, fatty acids, stabilisers, thickeners, colourants, flavourants, and the like. It is within the skills of the skilled person to select such further food ingredients depending on the desired end product.

[0030] In yet another aspect, the invention is concerned with a mixed bacterial culture comprising *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, and a strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in an aqueous medium comprising 0.25-10% (w/w) soy protein. The strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in such medium may advantageously be a *Lactobacillus plantarum* strain, preferably selected from the group of *Lactobacillus plantarum* B1839, B2484, and B2485. More preferably, said strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in such medium is *Lactobacillus plantarum* B1839. In a further embodiment, said *Lactobacillus delbrueckii* *bulgaricus* is selected from the group of *Lactobacillus delbrueckii* *bulgaricus* ATCC-BAA365, ATCC-11842 and Ib. The culture may be a yogurt starter culture, further comprising *S. thermophilus*. Alternatively, said strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in an aqueous medium comprising 0.25-10% (w/w) soy protein may be added to the aqueous medium as (part of) an adjunct culture in addition to a starter culture comprising *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, and *S. thermophilus*.

[0031] The invention also relates to the use of *Lactobacillus plantarum* B1839, B2484, and/or B2485, to stimulate growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, preferably *Lactobacillus delbrueckii* *bulgaricus* ATCC-BAA365, ATCC-11842 or Ib, preferably ATCC-BAA365, in an aqueous medium comprising 0.25-10% (w/w) soy protein.

[0032] The invention further pertains to the use of *Lactobacillus plantarum* B1839, B2484 and/or B2485, preferably B1839, in mixed culture fermentation of an aqueous medium comprising 0.25-10% (w/w) soy protein.

[0033] In yet another aspect, the present invention provides for a method for identifying a lactic acid bacterial strain capable of stimulating growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in a medium, said method comprising the steps of: i) growing a monoculture of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in said medium; ii) growing a mixed culture of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, and said lactic acid bacterial strain in said medium; iii) comparing the growth of *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in said monoculture with that in said mixed culture; and iv) selecting the lactic acid bacterial strain which increases colony forming units *Lactobacillus delbrueckii*, preferably *Lactobacillus delbrueckii* subsp. *bulgaricus*, in said medium. Said medium preferably is a sucrose-containing medium, more preferably an aqueous medium comprising 0.25-10% (w/w) soy protein, such as soy milk.

[0034] It will also be clear that the description, drawings and examples are included merely to illustrate some embodiments of the invention, and not to limit the scope of protection. Starting from this disclosure, many more embodiments will be evident to a skilled person, which are within the scope

of protection and the essence of this invention and which are obvious combinations of prior art techniques and the disclosure of this patent.

### DRAWINGS

**[0035]** The invention is illustrated using FIG. 1, in which the effects of mixed culture growth on outgrowth of *L. plantarum* are compared to outgrowth in the corresponding monocultures, assessed for 41 individual *L. plantarum* strains. A. Milk fermentations, B. Soymilk fermentations. NIZO strain numbers and log CFU values for growth in monocultures are presented on the X-axis and the difference in outgrowth in the presence of *S. thermophilus*, *L. bulgaricus* or both is represented on the Y-axis (as difference in log<sub>10</sub> CFU values). Gray; mixed cultures with *S. thermophilus*, White; mixed cultures with *L. bulgaricus*, Dark grey; mixed cultures with both *S. thermophilus* and *L. bulgaricus*.

### EXAMPLES

#### Materials and Methods

##### Screening Approach

**[0036]** Mixed culture growth and acidification properties were studied in milk and in soy milk which were prepared by UHT treatment of reconstituted skimmed milk powder (Friesland Foods, Leeuwarden, the Netherlands) and Bene-soy soy milk powder (Devansoy, Carroll, Iowa, USA). *Lactobacillus delbrueckii* subsp. *bulgaricus* strain ATCC-BAA365 and 41 *Lactobacillus plantarum* strains (Table 1) were cultured in MRS at 42 and 37° C., respectively, and *Streptococcus thermophilus* strain CNRZ1066 was cultured in GM17 at 42° C. Single and mixed cultures for fermentation of milk and soymilk were prepared in duplicate by diluting cultures of each strain to an O.D. (600 nm) of ~1 and subsequently transferring 1% of the diluted culture to 10-ml polystyrene screw cap tubes containing 5-ml of freshly prepared substrate. Milk and soy milk cultures were incubated at 37° C. for 24 h and then transferred to fresh, preheated substrate using an inoculation needle. Before each transfer the cultures were mixed thoroughly for 15 min in an Infors Microtron II incubator (Infors AG, Switzerland). After 4 rounds of fermentation in milk or soy milk the cultures were examined for outgrowth of individual strains and end pH.

TABLE 1

The <i>Lactobacillus plantarum</i> strains.		
Strain	Source	Alternative name
B1836	human D/S	WCFS1
B2896	vegetables	ATCC14917
B2029	Dairy	MLC43
B2457	fermented meat	CHEO3
B2484	fermented meat	NCTH19-1
B2485	fermented meat	NCTH19-2
B2494	fermented meat	NCTH27
B2535	fruit	LD2(2)
B2741	vegetables	NOS140
B2753	Sourdough	Q2
B2757	Sourdough	H4
B2766	Sourdough	H14
B2776	Dairy	CECT4645
B2801	vegetables	KOG18
B2802	Dairy	KOG24
B2806	vegetables	LMG9208
B2814	fruit	Lp95
B2830		
B2831	Silage	

TABLE 1-continued

The <i>Lactobacillus plantarum</i> strains.		
Strain	Source	Alternative name
B2855	fermented meat	N58
B2877	fermented meat	X17
B2889	fruit	LAC7
B2891	vegetables	LD3
B2897	vegetables	DKO22
B1837	Human C/F	299
B1838	Human C/F	CIP104440
B1839	vegetables	SF2A35B
B1840	vegetables	NCIMB12120
B2256	Human C/F	CIP104441
B2257	Human C/F	CIP104450
B2258	Human U	CIP104451
B2259	Human D/S	CIP104452
B2260	Human C/F	299v, DSM9843
B2261	Silage	NC8
B2262	Silage	LM3
B2263	Silage	LP80
B2264	Silage	LP85-2
B2726	Silage	ATCC8014(2)
B3894	vegetables	NCDO1193
B3892	Human D/S	CIP102359
B3893	Human D/S	CIP104448
B3400	Dairy	LMG18021

#### Quantification of Growth; HT Plating

**[0037]** Colony forming units (CFU) were determined using a plating protocol based on the method described recently by Sieuwerts et al (LAM paper). Culture dilutions were prepared in TY broth using a multichannel pipette and were plated by pipetting 20 µl of each dilution on TY agar in 12 cm squared Petri dishes. *L. plantarum* plates contained 1% galactose and were incubated at 30° C. for 24-36 h. *S. thermophilus* was plated on TY agar of which the pH was adjusted to 7.5, and glucose (1%) was added as substrate. *L. delbrueckii* subsp. *bulgaricus* was plated on TY agar containing 1% mannose. For the (moderately thermophilic) yoghurt species the plates were first incubated at 37° C. for 4 h and then the temperature was increased to 45° C. *L. delbrueckii* subsp. *bulgaricus* plates were incubated under anoxic conditions in jars containing a mixture of nitrogen (85%), carbon dioxide (10%) and hydrogen (5%). CFU counts were determined after 24-36 h of incubation. End pH of the cultures was determined using hydroplates in a Tecan Safire II spectrophotometer (Tecan Austria GmbH, Austria).

#### Population Dynamics

**[0038]** Population dynamics were studied in batch fermentors containing 1 L substrate from which samples were taken every 1.5 h. CFU's were determined according to the protocol described above and in addition samples were prepared for quantification by real-time quantitative PCR by mixing 0.1 ml sample with 0.9 ml TE. After mixing the samples, cell pellets were collected by centrifugation at 14,000×g and stored at -20° C. Primers for quantification of individual species based on 16S ribosomal DNA sequences were adapted from literature or designed using Amplicon and verified using Netprimer and Primer Express (Table 2). DNA was isolated using the Norgen Milk Bacterial DNA Isolation Kit. The amount of starting material was decreased to 0.1 ml and



the digestion period increased to 1 h. Quantification was performed on an Applied Biosystems 7500 Fast detection system.

TABLE 2

Primers used for quantification of individual species based on 16S ribosomal DNA sequences. The DNA sequences of the primers used are underlined, whereas the DNA sequence of the entire amplified sequence is depicted.
<i>L. delbrueckii</i> bulgaricus strain ATCC-BAA-365 qPCR primer set Ldf4-Ldr4 <u>CAAGTTTGAAAGGCGGCGTAAGCTGTCACTTTAGGATGAGCCCGC</u> <u>GCGCATTAGCTAGTTGGTGGGGTAAAGGCTACCAAGGCAATGATG</u> <u>CGTAGCCGAGTTGAGAGACTGATCGGCCA</u>
<i>L. plantarum</i> WCFS1 qPCR primer set Lp2f-Lp2r <u>TGATCCTGGCTCAGGACGAA</u> CGCTGGCGCGTGCCCTAATACATGCA <u>AGTCGAACGAACCTCTGGTATTGATTGGTGCTTGCA</u>
<i>S. thermophilus</i> CNRZ1066 primer set Tf2-Tr2 <u>AGGTAACCTGCCTTGTAGCGGGGATACTATTGGAAACGATAGCT</u> <u>AATACCGCATAACATGGATGACACATGTCATTTATTTGAAAGGGG</u> <u>CAATTGCTCCACTACAAGATGGACCTGCGTTGTAT</u>

## Results

### Combinatorial Screening Approach

**[0039]** Each of 41 *L. plantarum* strains was screened for the ability to ferment milk and soy milk in mixed culture combinations with *S. thermophilus* and *L. delbrueckii* subsp. *bulgaricus* and compared to growth in mono-cultures. The screening approach was developed to meet several criteria. To allow adaptation of each species to the fermentation conditions and the presence of one or more of the other species each culture was transferred 3 times to fresh substrate. This also increased the chance of selecting individuals with properties of interest since it enables fast growing strains to outcompete other strains for which no or only minor interdependencies exist. The fermentation temperature was set at 37° C. to allow good growth of each species, and before transferring to fresh substrate each culture was mixed thoroughly. To minimize the chance of (cross-)contamination, cultures were transferred using inoculation needles. To assess outgrowth of each species after 4 rounds of culturing in milk or soymilk a previously developed plating protocol (Sieuwerts, S., F. A. de Bok, E. Mols, W. M. de Vos, and J. E. van Hylckama Vlieg. 2008. A simple and fast method for determining colony forming units. Lett Appl Microbiol 47:275-8) was optimized for each of the 3 species of interest. Culture dilutions were prepared in TY broth using a multichannel pipette and 20 µl of each dilution was spotted using a multichannel pipette on 12 cm squared petridishes filled with TY agar, 6 spots per row and using only the odd or even tips of the pipette in order to use the available space optimally. TY agar was selected for plating because all 3 species and 41 *L. plantarum* strains produced CFU's on it which were comparable to growth on MRS agar (for *L. delbrueckii* and *L. plantarum*) or GM17 agar with 1% glucose (for *S. thermophilus*). *L. delbrueckii* and *S. thermophilus* plates were incubated at 45° C. to exclude growth of *L. plantarum*. However, because this resulted in significantly less CFU's of these species when compared to incubation at 37° C., the plates were first incubated at 37° C. for 4 h and then the temperature was increased to 45° C. Growth of *L. delbrueckii* on *S. thermophilus* plates was excluded by increas-

ing the pH of the agar to 7.5, at which growth of *S. thermophilus* was not inhibited. Mannose was used as substrate for *L. delbrueckii* since API tests showed that *S. thermophilus* was not able to grow on it. Formation of colonies by *L. delbrueckii* was optimal under anaerobic conditions. *L. plantarum* plates contained galactose as selective substrate and these plates were incubated at 30° C. Culture dilutions were prepared in TY broth using a multichannel pipette and 20 µl of each dilution was spotted using a multichannel pipette on 12 cm squared Petri dishes filled with TY agar. For each culture dilutions 10<sup>-1</sup> to 10<sup>-5</sup> were plated, by which the lower detection limit was set at 500 CFU's per ml. Agar plates, on which in total 36 culture dilutions were plated, were air dried and then incubated. End pH values of the cultures were determined using hydroplates, which were calibrated by including buffers within the range of pH 4 to 9.

### Screening of Mixed Cultures in Milk

**[0040]** A collection of 41 *L. plantarum* strains for which selection was based on differences in isolation source, geographic origin, and phylogenetic distance was screened for outgrowth in mono- and mixed culture milk fermentations with *S. thermophilus* and *L. delbrueckii* subsp. *bulgaricus*. After 4 rounds of culturing in milk viable cell counts in monocultures of 5 *L. plantarum* strains were below the detection limit of 500 CFU/ml, while most of the other strains produced at least more than 10<sup>6</sup> CFU/ml under these conditions (FIG. 1). Although the CFU levels of 17 strains was more than 10<sup>7</sup>, none of the *L. plantarum* mono cultures in milk exceeded 10<sup>8</sup> CFU/ml. None of the strains was thereby able to lower the pH of the milk below 6.3 which is only 0.25 less than the average pH of the milk controls. With one exception none of the *L. plantarum* strains produced more CFU when cultured together with either *S. thermophilus* or *L. delbrueckii* in milk. However, 21 strains produced more CFU when both *S. thermophilus* and *L. bulgaricus* were present, and nearly all of these strains showed poor growth in monocultures (FIG. 1). The presence of *L. plantarum* did not have a significant effect on outgrowth of *S. thermophilus* or *L. bulgaricus* in milk.

### Screening of Mixed Cultures in Soymilk

**[0041]** Screening for outgrowth of mono- and mixed cultures was also performed on soymilk using the same species, strains and combinations as described above for milk. With 1 exception all *L. plantarum* strains and *S. thermophilus* were able to ferment soymilk in monoculture. After 4 rounds of fermentation in soymilk *L. bulgaricus* was also still detected but did not exceed 10<sup>6</sup> CFU/ml in any of the 5 monoculture controls (average; 3.1±1.7·10<sup>5</sup>) and thereby did not affect the pH. Outgrowth of several *L. plantarum* strains was reduced up to 2 log scales by the presence of *S. thermophilus*, while no substantial effect of *L. plantarum* on outgrowth of *S. thermophilus* was observed. Apart from a small stimulation of *L. plantarum* strain B1839 by *L. bulgaricus*, no substantial positive effects of *L. bulgaricus*, *S. thermophilus* or both on outgrowth on any of the *L. plantarum* strains was observed. The CFU levels of *L. bulgaricus* had dropped below the detection limit (500 CFU/ml) in nearly all 2- and 3-species combinations. However, in some combinations *L. bulgaricus* was still present and in the 2- and 3-species combinations with *L. plantarum* B1839, as well as in 3-species combinations with *L. plantarum* strains B2484 en B2485, its CFU levels were

10-100 times higher when compared to the corresponding cultures without *L. plantarum*. The levels in the co-culture with *L. plantarum* strain B2776 were comparable to the levels in *L. bulgaricus* monocultures, consistent with the inability of this strain to ferment soymilk.

#### Population Dynamics of Mono- and Mixed Culture Milk and Soymilk Fermentations

[0042] Reproducibility of the screening experiment was assessed by repeating the screening protocol for several mono- and mixed cultures of interest with *L. plantarum* strains B1836 (WCFS1), B2485 and B1839 in soymilk and strains B2258 and B3400 in milk. The CFU-values for soymilk were comparable to the initial screening experiment, although *L. bulgaricus* produced 20 times more CFU in the mixed culture with strain B2485. For milk the results were comparable for strain B2258, but for outgrowth of B3400 in the 3-species mixed culture the CFU values were over 10-fold less. Strains B1839 and B3400 were selected for studying the population dynamics of soymilk and milk fermentations, respectively. The screening protocol was repeated for each mono- and mixed culture, but the 4<sup>th</sup> fermentation was performed in batch (stirred) fermentors containing 1 L substrate from which samples were taken every 1.5 h. Growth of *L. plantarum* B3400 in milk under these conditions appeared to continue throughout the fermentation period with an intermediary lag phase between 4 and 12 h, and this pattern was similar for each of the 4 mixed culture combinations. Throughout the fermentation CFU values of B3400 for the 3-species fermentation were approximately 100 fold higher than the monoculture and between 10- and 1000-fold higher than each of the 2-species fermentations. The dynamics of *S.*

*thermophilus* were nearly identical for the 4 cultures in which this species was present, and was characterized by 2 logarithmic growth phases. The dynamics of *L. bulgaricus* monoculture was nearly identical to the mixed culture with *L. plantarum*, while the dynamics of this species in the co culture with *S. thermophilus* was very similar to the 3-species milk fermentation.

[0043] In soymilk *L. plantarum* strain B1839 grew exponentially in the first 10-12 h of the fermentation irrespective of the presence of one or both of the other species, and in none of the cultures substantial increases in CFU were observed for the final 12 h of the fermentations. In the mixed culture with *L. bulgaricus* the CFU levels of B1839 were approximately 10-fold higher than for each of the other cultures throughout the fermentation period. In the mono-culture and the mixed culture with both yoghurt species CFU levels still increased after 10 h while the level in the culture with *S. thermophilus* remained fairly constant after 10 h of fermentation. The behavior of *S. thermophilus* was nearly identical in each of the soymilk cultures with an exponential growth phase in the first 6 h of fermentation and no substantial increase in CFU levels for the remainder of the fermentation period. The CFU level of *L. bulgaricus* in the mono-culture did not exceed the detection limit, while its dynamics in the mixed cultures with *L. bulgaricus* and both other species were comparable. A substantial positive effect of *L. plantarum* on *L. bulgaricus* was observed, in particular when *S. thermophilus* was not present. The experiments were repeated with 2 different *L. delbrueckii* subsp. *bulgaricus* strains, i.e., ATCC-11842 and Ib, with comparable results. Thus, it is concluded that *L. plantarum* B1839 has a positive effect on outgrowth of *L. delbrueckii* subsp. *bulgaricus*.

---

#### SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 9

<210> SEQ ID NO 1  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: qPCR primer Ldf4

<400> SEQUENCE: 1

caagtttgaa aggcggcgta

20

<210> SEQ ID NO 2  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: qPCR primer Ldr4

<400> SEQUENCE: 2

gttgagagac tgatcggccca

20

<210> SEQ ID NO 3  
<211> LENGTH: 20  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:

-continued

---

```

<223> OTHER INFORMATION: qPCR primer Lp2f

<400> SEQUENCE: 3

tgatcctggc tcaggacgaa                20

<210> SEQ ID NO 4
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: qPCR primer Lp2r

<400> SEQUENCE: 4

tgggtattgat tgggtgcttgc a            21

<210> SEQ ID NO 5
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: qPCR primer Tf2

<400> SEQUENCE: 5

aggtaacctg ccttgtagcg g              21

<210> SEQ ID NO 6
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: qPCR primer Tr2

<400> SEQUENCE: 6

caagatggac ctgcgttgta t              21

<210> SEQ ID NO 7
<211> LENGTH: 121
<212> TYPE: DNA
<213> ORGANISM: Lactobacillus delbrueckii bulgaricus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(121)
<223> OTHER INFORMATION: strain ATCC-BAA-365 amplified fragment

<400> SEQUENCE: 7

caagtttgaa aggcggcgta agctgtcact ttaggatgag cccgcggcgc attagctagt    60
tgggtgggta aaggcctacc aaggcaatga tgcgtagccg agttgagaga ctgacggcc      120
a                                          121

<210> SEQ ID NO 8
<211> LENGTH: 81
<212> TYPE: DNA
<213> ORGANISM: Lactobacillus plantarum
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(81)
<223> OTHER INFORMATION: strain WCFS1 amplified fragment

<400> SEQUENCE: 8

tgatcctggc tcaggacgaa cgctggcggc gtcctaata catgcaagtc gaacgaactc    60
tgggtattgat tgggtgcttgc a                                          81

```

-continued

```

<210> SEQ ID NO 9
<211> LENGTH: 127
<212> TYPE: DNA
<213> ORGANISM: Streptococcus thermophilus
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(127)
<223> OTHER INFORMATION: strain CNRZ1066 amplified fragment

<400> SEQUENCE: 9

aggtaacctg cctgtgtagcg ggggataact attggaaacg atagctaata ccgcataaca      60
atggatgaca catgtcattt atttgaaagg ggcaattgct ccactacaag atggacctgc      120
gttgatat                                     127

```

1. A method for preparing a fermented soy product, said method comprising: fermenting an aqueous medium comprising 0.25-10% (w/w) soy protein with a mixed bacterial culture comprising, in addition to *Lactobacillus delbrueckii*, a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

2. The method according to claim 1, wherein said strain capable of stimulating growth of *Lactobacillus delbrueckii* is a *Lactobacillus plantarum* strain, optionally comprising *Lactobacillus plantarum* B1839, B2484 and/or B2485.

3. The method according to claim 1, wherein said *Lactobacillus delbrueckii* is *Lactobacillus delbrueckii* subsp. *bulgaricus*, optionally comprising *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC-BAA365, ATCC-11842 or Ib.

4. The method according to claim 1, wherein the fermented soy product comprises soy yogurt.

5. A fermented soy product comprising *Lactobacillus delbrueckii* and a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

6. The fermented soy product according to claim 5, wherein said strain capable of stimulating growth of *Lactobacillus delbrueckii* is a *Lactobacillus plantarum* strain, optionally comprising *Lactobacillus plantarum* B1839, B2484, and/or B2485.

7. The fermented soy product according to claim 5, wherein said *Lactobacillus delbrueckii* is *Lactobacillus delbrueckii* subsp. *bulgaricus*, optionally comprising *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC-BAA365, ATCC-11842 or Ib.

8. The fermented soy product according to claim 5, wherein said soy product comprises an aqueous soy product comprising 0.25-10% (w/w) soy protein.

9. The fermented soy product according to claim 5, wherein said soy product comprises soy yogurt.

10. A culture comprising *Lactobacillus delbrueckii* and a strain capable of stimulating growth of *Lactobacillus delbrueckii*.

11. The culture according to claim 10, wherein said strain capable of stimulating growth of *Lactobacillus delbrueckii* is

a *Lactobacillus plantarum* strain, optionally comprising *Lactobacillus plantarum* B1839, B2484 and/or B2485.

12. The culture according to claim 10, wherein said *Lactobacillus delbrueckii* *bulgaricus* is *Lactobacillus delbrueckii* subsp. *bulgaricus*, optionally comprising *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC-BAA365, ATCC-11842 or Ib.

13. A *Lactobacillus plantarum* B1839 capable of being used to stimulate growth of *Lactobacillus delbrueckii*, optionally comprising *Lactobacillus delbrueckii* subsp. *bulgaricus*, optionally comprising *Lactobacillus delbrueckii* subsp. *bulgaricus* ATCC-BAA365, ATCC-11842 or Ib., in an aqueous medium comprising 0.25-10% (w/w) soy protein.

14. A *Lactobacillus plantarum* B1839, B2484 and/or B2485, optionally comprising B 1839, capable of being used in mixed culture fermentation of an aqueous medium comprising 0.25-10% (w/w) soy protein.

15. A method for identifying a lactic acid bacterial strain capable of stimulating growth of *Lactobacillus delbrueckii* in a medium, said method comprising:

- a) growing a monoculture of *Lactobacillus delbrueckii* in said medium;
- b) growing a mixed culture of *Lactobacillus delbrueckii* and said lactic acid bacterial strain in said medium;
- c) comparing growth of *Lactobacillus delbrueckii* in said monoculture with growth in said mixed culture; and
- d) selecting a lactic acid bacterial strain which increases colony forming units *Lactobacillus delbrueckii* in said medium.

16. The method according to claim 15, wherein said medium is a sucrose-containing medium, optionally comprising an aqueous medium comprising 0.25-10% (w/w) soy protein, optionally comprising soy milk.

\* \* \* \* \*